

---

# BI-ACTIV®

---

Alcoholic fermentation booster to be used in the case of sluggish or stuck fermentations.

*Qualified for the elaboration of products for direct human consumption in the field of the regulated use in Oenology.  
In accordance with the current EU regulation n° 2019/934.*

## SPECIFICATIONS

---

A specific formulation based on yeast cellular envelopes (yeast hulls), inert support elements (cellulose), and inactivated yeast for vinification, **BI-ACTIV®** provides:

- A physical support effect for the yeasts.
- Detoxification of the must and the wine.

## OENOLOGICAL APPLICATIONS

---

In the event of slow alcoholic fermentation, **BI-ACTIV®** enables the grape must to be detoxified, provides survival factors for the yeasts and enables fermentation to be completed.

In the event of stuck fermentation, **BI-ACTIV®** detoxifies the wine and prepares it for the new inoculation.

**BI-ACTIV®** can be used in the vinification of highly clarified must and immediately provides support elements and survival factors, which are essential for membrane stress-resistance.

**BI-ACTIV®** does not provide assimilable nitrogen.

**BI-ACTIV®** can be used on all types of must or wine, white, rosé or red.

## SCIENTIFIC RESULTS

---

The inert elements allow a support effect for the yeast, essential in the case of highly clarified must. The yeast walls have a high adsorption capacity for medium-chain fatty acids (C6, C8, C10) (Lafon-Lafourcade *et al.*, 1984), which act as fermentation inhibitors (Salmon *et al.*, 1993). Finally, inactivated yeasts provide survival factors (long chain fatty acids and sterols).

**BI-ACTIV®** also has a positive effect on malo-lactic fermentation.

## PHYSICAL CHARACTERISTICS

---

Aspect ..... powder      Apparent density (tight packed) (g/L) ..... ≈ 500  
Color ..... white



**LAFFORT**

*l'œnologie par nature*

## CHEMICAL AND MICROBIOLOGICAL ANALYSIS

Total nitrogen (%) ..... ≈ 1	Lead (ppm) ..... < 2
<i>E. coli</i> (/25 g) ..... none	Arsenic (ppm) ..... < 3
<i>Staphylococcus</i> (/g) ..... none	Mercury (ppm) ..... < 1
<i>Salmonella</i> (/25 g) ..... none	Cadmium (ppm) ..... < 1

## PROTOCOL FOR USE

### OENOLOGICAL CONDITIONS

- In the event of a stuck fermentation, adjust the SO<sub>2</sub> (20 to 30 ppm), rack and then add the **BI-ACTIV**<sup>®</sup>. *Please refer to our restarting stuck fermentations protocol.*

### DOSAGE

- In the event of slow fermentation (white, rosé, red) and low turbidity: 30 g/hL (300 ppm).
- In the event of treatment for a stuck fermentation: 60 g/hL (600 ppm) for red, 30 g/hL (300 ppm) for white/rosé or if used in conjunction with **TURBICEL**.

Maximum legal dose (EU): 303 g/hL (3030 ppm).

### IMPLEMENTATION

In order to allow for optimal expansion of the support elements, leave the product to aerate for 10 minutes before use. The product must be used within 1 hour of opening. Do not use opened bags. Use a clean, inert container. Dissolve the total quantity of **BI-ACTIV**<sup>®</sup> to be added in 10 times its weight in must or wine. Mix well, then incorporate directly into the tank during a pump-over.

### STORAGE RECOMMENDATION

- Store above ground level in a dry area not liable to impart odours. Ensuring stock is kept at a moderate temperature, in its original, unopened packaging.
- Optimal date of use: 3 years.

### PACKAGING

1 kg bag. 10 kg box.

