

# QUERTANIN® PLUS

Stave wood quality ellagic tannins, extracted from american oak heartwood. For post-vinification use in red, rosé and white wine.

*Qualified for the elaboration of products for direct human consumption in the field of the regulated use in oenology.  
In accordance with the current regulation (EU) 2019/934.*


## SPECIFICATIONS AND OENOLOGICAL APPLICATIONS

**QUERTANIN® PLUS** is a preparation of ellagic tannins used in vinification or during ageing, with antioxidant and antioxidase properties (inhibition of laccase) and high reactivity with proteins (facilitates clarification).

- Regulates redox phenomena during barrel ageing or micro-oxygenation.
- Provides older barrels with an anti-oxidant medium rich in ellagitannins, identical to that of new barrels.

## EXPERIMENTAL RESULT

- Tannin content extracted from the oak is lower in used barrels. The ellagitannin protective effect is decreased and the wine becomes subject to premature oxidation. Adding **QUERTANIN® PLUS** permits the re-creation of the buffer qualities provided by tannins extracted from new barrels, and protects the wine from oxidation phenomena.



## PHYSICAL CHARACTERISTICS

Aspect ..... granulated  
Solubility ..... complete

Colour ..... dark brown



**LAFFORT**

*l'œnologie par nature*

## CHEMICAL ANALYSIS

Tannins (%) .....	> 65	Arsenic (ppm) .....	< 3
Humidity (%) .....	< 10	Iron (ppm) .....	< 50
Ash (%) .....	< 5	Lead (ppm) .....	< 2
Insoluble substances (%) .....	< 5	Mercury (ppm) .....	< 0.5
		Cadmium (ppm) .....	< 0.5

## PROTOCOL FOR USE

### DOSAGE

*It is specified in the Oenological Codex that tannins "must not alter the recognised organoleptic characteristics of the wines". The dosage rates will therefore vary in function of the wine matrix, and shall be determined after trials.*

- Clarification additive: 10 – 30 g/hL (100 - 300 ppm).
- Inhibition of laccase activity: 30 – 80 g/hL (300 - 800 ppm).
- Prevention of oxidation in white and rosé wines: 2 - 5 g/hL (20 - 50 ppm).
- Red wine ageing: 5 - 20 g/hL (50 - 200 ppm).

### IMPLEMENTATION

It is recommended to add **QUERTANIN® PLUS** at least two weeks before bottling.

*For white wines stabilised with CMC (tartaric stabilisation), the addition of tannins towards the end of ageing may affect the colloidal stability; it is recommended to perform a protein stability test again systematically after addition.*

## STORAGE RECOMMENDATION

- Store above ground level in a dry area not liable to impart odours. Ensuring stock is kept at a moderate temperature, in its original, unopened packaging.
- Optimal date of use: 5 years.

## PACKAGING

500 g bag - 10 kg box.

