SUPERSTART® Blanc

Yeast preparate to use for in ADY rehydration phase for a strong fermentation finish. Patent FR 2736651. Specific formulation for white and rosé wines conditions.

Qualified for the elaboration of products for direct human consumption in the field of the regulated use in Oenology.

In accordance with the current EU regulation n° 2019/934.

SPECIFICATIONS AND OENOLOGICAL APPLICATIONS

A specific preparation of yeast origin naturally rich in vitamins, minerals, long chain fatty acids and sterols. The growth factors of **SUPERSTART® BLANC** help regulate the formation of undesired compounds and the yeast growth and are cofactors of membrane transports. The survival factors are implied in maintaining the conformation and membrane resistance.


Thanks to its specific formulation for white and rosé wines conditions, **SUPERSTART® BLANC** optimises the potential of the fermenting yeast to produce and release aromatic compounds. In addition it guarantees a better resistance to alcohol and a strong fermentation finish.

SUPERSTART® BLANC increases yeast viability and the general metabolism of the yeast and therefore:

- Increases the yeasts' resistance to difficult conditions (low turbidity, anaerobic winemaking, high alcohol concentrations).
- Optimises the revelation and production of aromatic compounds by yeasts.
- · Prevents the excessive production of volatile acidity.
- Reinforces the efficiency of «pied de cuve».


EXPERIMENTAL RESULTS

· Better cell viability



The evolution of the cell population by **ZYMAFLORE® X16** rehydrated with and without **SUPERSTART® BLANC**. Fermentations are done on a 200 g/L of sugar must and a level of available nitrogen < 140 mg/L.

· Improved yeast aromatic performance



Sauvignon Blanc, France, TAV 12.5% vol. Differences perceived during tasting.

By improving overall assimilation of must compounds, SUPERSTART® BLANC optimises the metabolism of the yeast to produce more aromatic wines.



PHYSICAL CHARACTERISTICS

Aspect	powder	Colour	beig	2e

CHEMICAL & MICROBIOLOGICAL ANALYSIS

Humidity (%)< 7	Lead (ppm)
Total nitrogen (%)< 11	Arsenic (ppm)
E. coli (/g)none	Mercury (ppm)
Staphylococcus (/g)none	Cadmium (ppm)
Salmonella (/25 g)none	

PROTOCOL FOR USE

OENOLOGICAL CONDITIONS

To be added to yeast rehydration water. Do not use directly in the tank (the elements provided by **SUPERSTART® BLANC** would be absorbed by the indigenous flora or would be bound by certain components).

DOSAGE

20 - 30 g/hL (200 - 300 ppm) of must to be fermented, in the active dry yeast rehydration water.

......< 2
.....< 3
.....< 1
.....< 1

SUPERSTART® BLANC is recomended for the initial yeast inoculation and for the restart culture in the event of a stuck fermentation. (in this case, refer to our fermentation restart protocol).

IMPLEMENTATION

Do not use open bags.

Use a clean, inert container. Dissolve the total quantity of **SUPERSTART® BLANC** needed for the fermentation tank in 20 times its weight in water at 37°C (98.6°F). Mix well, then incorporate the active dry yeast. Follow the protocol for standard active dry yeast rehydration (refer to the yeast packaging).

STORAGE RECOMMENDATION

- Store above ground level in a dry area not liable to impart odours. Ensuring stock is kept at a moderate temperature, in its original, unopened packaging.
- · Optimal date of use: 3 years.

PACKAGING

1 kg bag - 10 kg box.

5 kg bag - 10 kg box.

