

Flotation Protocol with VEGECOLL®

Many factors influence the flotation process and hence its success. The parameters of the protocol have been specially adapted for an easy flotation. Do not hesitate to contact **LAFFORT®**'s team before your flotation trials in order to explore the potentially inhibiting parameters and find the appropriate solutions.

1 PREPARING OF THE JUICE


- Clarification by flotation involves migration of the particles of the must to the surface of the tank. This migration is prevented in the presence of pectins. The addition of pectolytic enzyme directly after grape pressing is necessary to accelerate the process.
 - **LAFASE® XL FLOT** - 1 - 4 mL/hL.
 - **LAFAZYM® 600XL^{ICE}** (allows complete depectinization at low temperatures) 1 - 2 mL/hL.
- In the case of must particularly difficult to clarify (variety, maturity, ...) or to accelerate the depectinization the use of **LAFASE® BOOST** at 1 mL / hL is recommended.
- Check the completion of the depectinization before starting the flotation. Use our **PECTIN TEST**, easy and fast.



2 CONNECTING THE FLOTATION PUMP

- For an easy flotation, the filling of the tank should not exceed 85 to 90% of the total volume.
- The temperature of the must should be between 15 and 18°C (59-65°F). The colder the must, the higher the viscosity, the more difficult the flotation process.
- Connect the pump inflow to racking valve, and the pump outflow to lees valve.
- For the best results, total hose length should not exceed 3 m (inflow and outflow).
- Make sure all the air is out of the saturation column before closing the tap.


MUST FOR CLARIFICATION



3 ADDING THE VEGECOLL®

- Start the pump without gas injection.
- Check that the saturation pressure is between 2 and 3 bar (the size of the tank does not matter). Prepare **VEGECOLL®** in a clean, inert container following **LAFFORT®** recommendations.
- The recommended dose of **VEGECOLL®** is 20 - 100 ppm. (the dose can be adjusted according to the characteristics of the must).
- Place into the **VEGECOLL®** preparation the pipe dedicated to the venturi section provided on the flotation system.
- Inject **VEGECOLL®** as slowly as possible.
- Mix the tank for 20 to 25 minutes at a saturation pressure of 2 to 3 bar, without addition of gas.

MUST FOR CLARIFICATION




Flotation Protocol with VEGECOLL®

4 STARTING THE FLOTATION PROCESS


- Once the tank is homogenised, open the gas injection valve.
- The nitrogen inlet pressure should be between 5 and 7 bar.
- The gas flow rate must be between 25-60 L/min (depending on flotation setup). The saturation pressure must be adjusted to 5 bar.
- Check the quality of the flotation. To do this, take a sample at the tap of the saturation column.
- Remember to readjust the saturation pressure between 5 and 7 bar after sampling.
- The circulation time for flotation is between 60-150 min. Depending on the volume of the tank.
- Pump the equivalent of 1 to 2 volumes of the tank - 1.5 times is usually enough.

MUST FOR CLARIFICATION



5 COMPLETION OF FLOTATION PROCESS AND WAITING TIME

- Once the flotation process is complete, stop the pump.
- Close the gas.
- Close all the valves in the tank.
- Leave the tank for 60 to 120 minutes so that lees can rise to the surface.
- **Do not leave the tank longer than 240 minutes. Gravitational force can cause lees separation and resuspension of the lees if the waiting time is too long.**
- Check the turbidity of the clarified batch.



Find out more: Discover our *FLOTATION* video on our website, at **LAFFORT & YOU**.